





# KynoPop®



#### Name:

#### **KynoPop®**

# Properties (What):

A fine, dry highly water-soluble product.

KynoPop®	N	Р	К	Mg	S
Macro elements (g/kg)	143	87,5	42,3	-	-
Secondary elements (g/kg)	-	-	-	18	26

KynoPop®	Zn*	Cu*	В	Мо
Micro elements (mg/kg)	24000	6000	2000	400

<sup>\* =</sup> chelated

# Benefits (Why):

- Although there may be sufficient nutrients in the soil, there may be various reasons for maize seedlings to experience nutrient deficiencies, especially at seedling establishment.
- Limited root development during the seedling establishment stage, can impair nutrient uptake.
- Cold, wet conditions can often occur at planting. Such conditions can impair the uptake of nutrients and **KynoPop®** can lower this risk.
- Certain nutrients are more important than others during various growth stages. The product has been specifically developed to provide for the immediate needs of the young seedling.
- All the raw materials in KynoPop® are highly "plant available" and are
  "soft" products that should, at recommended rates, not damage the
  young roots of germinating seeds. The micro elements are chelated to
  enhance plant availability.

- KynoPop® enhances the effectiveness of the seedling and improves early root development.
- Stronger seedlings can better tolerate stress conditions (cold, heat, drought, herbicides, etc.), diseases and pests.

## Application (How):

 As a single soil application (10% solution) of KynoPop® with planting:

Row width	Application rate
225 cm	4 – 7 kg/ha
152 cm	5 – 8 kg/ha
91 cm	6 – 9 kg/ha
76 cm	7 – 10 kg/ha
52 cm	9 – 12 kg/ha

 If is to be applied at a later stage the above application rate (10% solution) can be reduced by 20%.

## Uses (Where):

- KynoPop® contains macro, secondary and micro elements in a ratio specifically beneficial to seedlings.
- Extremely suitable in cold and wet conditions during planting.
- Extremely suitable for sandy soils where nutritional deficiencies can handicap young seedlings and where wind damage can occur.