


# Cabrio® Duo

Reg. No:



**NET CONTENTS:**  
**1.0 LITRE**

**HARMFUL IF SWALLOWED**

<b>Composition</b>	<b>mass/volume</b>
Pyraclostrobin.....	40g/L
Dimethomorph.....	72g/L
Inert ingredients.....	888g/L

**Cabrio® Duo is an emulsifiable concentrate systemic fungicide for the control of frog-eye and alternaria in tobacco.**

**Registered by:**

**Pivotal Agro-Services (Pvt) Ltd.**

Agriculture House, Adylinn Road, Marlborough  
PO Box WGT 450, Westgate, Harare  
Telephone (263-4) 309800 / 18

**Manufactured by:**

**BASF SE Germany.**

®Registered trademark of BASF

**TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.**

## WARNINGS

1. Harmful if swallowed.
2. Irritating to the skin.
3. Highly toxic to aquatic organisms. May cause long term adverse effects to the aquatic environment.
4. Keep out of reach of children.

## PRECAUTIONS

1. KEEP AWAY FROM FOOD OR/AND BEVERAGES.
2. **DO NOT** eat, drink or smoke during use. Avoid contact with skin.
3. Wear protective clothing when handling or using this pesticide e.g. rubber gloves, face mask etc.
4. This material and its container must be disposed of as hazardous waste. **DO NOT** dispose of in the environment.
5. **DO NOT** contaminate water with the product or its container.
6. To protect aquatic organisms, leave a safety distance of about 5 meters between treated areas and surface water bodies.
7. Empty Container Disposal: TRIPLE RINSE empty containers in the following manner. Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds until the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinsings to the contents of the spray tank. Destroy the container by perforation, flattening and burying in an approved dump site. Never re-use for any other purpose.
8. Prevent contamination of food, feed, drinking water and eating utensils.

## FIRST AID TREATMENT

General advice:

Avoid contact with the skin, eyes and clothing. Take off all contaminated clothing immediately. If difficulties occur, seek medical attention. Show container, label and/or safety data sheet to physician.

If inhaled:	Keep the patient calm. Remove the patient to fresh air. Seek medical attention.
On skin contact:	After contact with skin, wash immediately with plenty of water and soap. If irritation develops, seek medical attention.
On contact with eyes:	Immediately wash affected eyes for at least 15 minutes under running water with eyelids held open. Consult an eye specialist.
On ingestion:	Immediately rinse mouth and then drink plenty of water. DO NOT induce vomiting. Seek medical attention. Never induce vomiting or give anything by mouth if the victim is unconscious or having convulsions.

Note to physician:

Treatment: Treat according to symptoms (decontamination, vital functions). No known specific antidote.

Symptoms of poisoning: eyes and skin irritation, skin dryness or cracking

## DIRECTIONS FOR USE

- **DO NOT** apply more than two applications of Cabrio® Duo in a row and more than 3 applications per year
- Apply enough volume of spray solution that will enable a complete and uniform wetting of the crop, but minimize dripping
- Cabrio® Duo should be applied preventively.
- It is recommended to use Cabrio® Duo during the critical periods of development of crop diseases listed.
- Shorter intervals between treatments are recommended in conditions with a high disease pressure, heavy rainfall or rapid growth.
- Use Cabrio® Duo as part of a spray program that includes other fungicides from different classes with different modes of action
- **DO NOT** apply Cabrio® Duo until the leaves are dry

### Preparation of the Mixture

Ensure that the spraying equipment is clean and properly calibrated.

1. Fill the spray tank half full with clean water.
2. Start the agitation system then add the required amount of Cabrio® Duo to the tank.
3. Top up the required volume of water, while agitating the solution.
4. **DO NOT** allow the mixture to sit overnight.
5. After the application, it is good practice to clean the spraying equipment.

**Compatibility:** When mixing Cabrio® Duo with other products, always perform preliminary testing on a limited number of plants before extending full treatment.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weeds against the remedy concerned as well as the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animals or for the lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of the conditions which could not have been foreseen in terms of registration. Consult the supplier in the event of any uncertainty.

### APPLICATION RATES

Crop	Diseases	Dose rate	Spray interval
Tobacco	Frog Eye ( <i>Cercospora nicotianae</i> )  Alternaria ( <i>Alternaria alternata</i> )	1.5- 2.5 l/ha in 200 litres of water  Use the higher rate under conditions of high disease pressure.	Begin spraying at 8 weeks after planting.  7-10 days